


- _____ After the mailing date of a final action under 37 C.F.R. §1.113, provided that this occurs prior to the issuance of a Notice of Allowance and provided that this I.D.S. is accompanied by either a certification as specified in 37 C.F.R. §1.97(e), a petition requesting consideration of the I.D.S., and the petition fee set forth in 37 C.F.R. §1.17(i)(1); and
- _____ After the mailing date of a Notice of Allowance under 37 C.F.R. §1.311, provided that this occurs prior to the issuance of a final action and provided that this I.D.S. is accompanied by either a certification as specified in 37 C.F.R. §1.97(e), a petition requesting consideration of the I.D.S., and the petition fee set forth in 37 C.F.R. §1.17(i)(1).

It should be understood that attention has been called to the references that have been deemed to be pertinent to the claimed present invention. In concluding what was pertinent, the criteria employed was considered most appropriate in light of the invention shown in the present application. However, the Examiner or others may deem some other criteria to be just as appropriate or more appropriate. Therefore, the Examiner is respectfully urged to review the listed references and to make the usual careful independent search for other prior art that may be pertinent.

Respectfully submitted,

July 15, 2002

Date



Paul D. Greeley
Reg. No. 31,019
Ohlandt, Greeley, Ruggiero & Perle, L.L.P.
One Landmark Square, 10th Floor
Stamford, CT 06901-2682
(203) 327-4500

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A24D 3/14, 3/16	A1	(11) International Publication Number: WO 96/00019 (43) International Publication Date: 4 January 1996 (04.01.96)
(21) International Application Number: PCT/GR94/00015 (22) International Filing Date: 27 June 1994 (27.06.94) (71)(72) Applicants and Inventors: STAVRIDIS, Ioannis [GR/GR]; 15 Chremonidou Street, GR-166 33 Athens (GR). DELICONSTANTINOS, George [GR/GR]; 40-44 Ilios Street, GR-115 27 Athens (GR). (74) Agent: LABROPOULOS, Socrates; 18 Akadimias Street, GR-106 71 Athens (GR).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: REMOVAL OF NOXIOUS OXIDANTS AND CARCINOGENIC VOLATILE NITROSOCOMPOUNDS FROM CIGARETTE SMOKE USING BIOLOGICAL SUBSTANCES (57) Abstract This invention refers to a method of withholding noxious compounds contained in cigarette smoke (NO, NOx, carcinogenic nitrosocompounds, free radicals, H ₂ O ₂ , CO, aldehydes, and trace elements) which were up to today insufficiently retained by conventional cigarette filters. The method described specifically refers to the enrichment of common convention filters with biological substances of the metal ions (Fe ²⁺ , Cu ²⁺ , Mg ²⁺) complexed with porphirin ring as well as Fe ²⁺ ions stereospecifically bound to protein molecules, either separately or in combinations. The enrichment of these conventional filters with the abovementioned biological substances alters neither the physical properties of the cigarette smoke (odor, taste and appearance) nor the physical properties of the filter itself.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Removal of noxious oxidants and carcinogenic volatile nitrosocompounds from cigarette smoke using biological substances.

5

The present invention establishes a methodology for withholding the noxious compounds, ie. nitrogen oxides, free radicals, aldehydes, hydrogen peroxide, carbon monoxide, trace elements and carcinogenic volatile nitrosocompounds from being inhaled during cigarette smoking, substances which until today are insufficiently retained by the use of conventional cigarette filters.

THEORETICAL BACKGROUND-LEVEL OF PREVIOUS TECHNOLOGY

15 A plethora of publications in international journals suggests that cigarette smoke is separated into two phases: a) a solid phase (tar); and b) a gas phase. This separation occurs with the use of a typical Cambridge-glass-fiber filter which withholds 99.9% of the particles which are greater in size than 0.1 μm . The tar of the cigarette contains dramatically high concentrations of very stable free radicals which can be classified into at least four different categories. Semiquinones in equilibrium with quinone and hydroxyquinones are considered to be free radicals with most interesting chemical properties. The quinone system reduces the molecular oxygen to form superoxide (O_2^-) which then upon spontaneous dismutation forms hydrogen peroxide (H_2O_2). In the gas phase, there are more than 10^{15} organic radicals per puff with half-lives of less than 20 1 second that are inhaled. It is paradoxical however that despite their minute half life these radicals can maintain high levels of activity for more than 10 minutes in the gas phase. In fact the concentration of these radicals is considerably increased as we approach the filter-end of the cigarette. An explanation for this paradox is to be found in the maintenance of a steady state situation; due to the ongoing production of free radicals (Pryor, W.A., Stone, K., Ann. N.Y. Acad. Sci. 686: 12-28, 1993).

30 Nitric oxide (NO) is the most important free radical in the gas phase of the cigarette smoke which, during smoking, participates in a sequence of reactions through which nitrogen dioxide, isoprene radicals, peroxy radicals and alkoxy radicals are

formed. Cigarette smoke also contains a considerable number of aldehydes which contribute to its damaging toxic effects. It has been shown that minute amounts of aldehydes extracted from the cigarette smoke cause both protein catabolism and oxidation of thiol groups of the plasma proteins. These properties attributed to the aldehydes are the result of the reactions between the carbonyl group of the aldehydes and the -SH and -NH₂ groups of the plasma proteins. For example, acrolein, from the cigarette smoke, reacts quickly with the -SH groups to form carbonyl compounds (Alving, K., Forhem, C., and Lundberg, J.M., Br. J. Pharmacol. 110: 739-746, 1993). In the tar of the cigarette smoke there are trace elements of, for example, iron, copper, manganese and cadmium which are implicated in many free radical producing reactions and lead to the formation of very active secondary radicals (e.g. peroxy radicals, alkoxy radicals, superoxide, cytotoxic aldehydes etc.). The introduction of the trace elements into the lung during cigarette smoking leads to a series of redox reactions both in lung fluids and alveolar macrophages which result in the formation of the very active hydroxyl radicals (OH[•]). These hydroxyl radicals are mainly formed in the presence of iron via the Fenton reaction. Copper can also form hydroxyl radicals by reacting with the hydrogen peroxide in the lung. Manganese, in low concentrations (10⁻⁷ M), stimulates the soluble guanylate cyclase of the endothelial cells of the lung causing the production of nitric oxide and superoxide through a positive feedback mechanism (Youn, Y.K., Lalonde, C., and Demling, R., Free Rad. Biol. Med. 12: 409-415, 1992). Carbon monoxide is produced during tobacco burning. A quantity of CO is retained in the lung even after exhaling, resulting in the stimulation of the soluble guanylate cyclase after its interaction with the heme moiety of the enzymes of the endothelial cells and other cells of the lung tissue. The increased levels of cyclic GMP within the cells coupled with a positive feed back mechanism increase the production of nitric oxide and superoxide (Watson, A., Joyce, H., Hopper, L., and Pride, N.B., Thorax 48: 119-124, 1993). NO gas which can be produced by numerous cell types, including the vascular endothelial cells and reticular endothelial cells, causes relaxation of the smooth muscle (Lowenstein, C.J., Dinerman, J.L., Snyder, S.H. Ann. Intern. Med. 120: 227-237, 1994). There are also exogenous sources of NO which are considered similarly responsible in causing damage to the blood vessels and other tissues. It is well established that secondary and tertiary amines can react with nitrite and other nitrosating agents to form N-nitrosoamines (Lowenstein, C.J., Dinerman, J.L., Snyder, S.H. Ann

Intern. Med. 120: 227-237, 1994). Since 1974 a number of studies have demonstrated that during harvesting, tobacco processing and smoking the alkaloids are nitrosated to tobacco specific N-nitrosamines (TSNA). Of the TSNA identified in tobacco and/or its smoke, N-nitrosonornicotine (NNN), 4-(methylnitrosoamino)-1-(3-pyridyl)-1-butanone (NNK) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) are strong animal carcinogens. NNN induces tumor of the lung in mice, tumors of the trachea in hamsters, and tumors of the nasal cavity and esophagus in rats. NNK induces tumors of the lung in mice, hamsters and rats, and also tumors of the liver, nasal cavity and pancreas in rats. Oral swabbing of a mixture of NNN and NNK elicits tumors in the oral cavity and lung of rats. The typical amount of both NNK and NNN in mainstream cigarette is 200 ng/cigarette. (Hecht, S.S., Spratt, T.E., and Trushin, N. Carcinogenesis, 9: 161-165, 1988).

Our present research, related to the effect of the cigarette smoke on lung tissue has revealed that NO reacts with superoxide to form the strong oxidant radical peroxynitrite (ONOO⁻) which causes secondary damaging reactions in key biomolecules. Both the metabolic and damaging effects of the NO in the cells were studied in our lab in vitro and in vivo experiments.

NO is oxidized, in the presence of oxygen, to nitrogen dioxide (NO₂). The rate of this oxidation depends upon the concentration of oxygen and the square of the NO concentration. Nitrogen dioxide is clearly cytotoxic and is transformed into nitrite and nitrate when in water solutions. Moreover NO forms complexes with trace elements and/or with metalloproteins, hemoglobin for example (Wink, D.A., Darbyshire, J.F., Nims, R.W., Saavedra, J.E., and Ford, P.E., Chem. Res. Toxicol. 6: 23-27, 1993).

NO that reacts with superoxide to form the noxious compound ONOO⁻ can justify certain types of superoxide toxicity. ONOO⁻ is unusually stable, taking into consideration its strong oxidative potential (+1.4 V). During its decomposition it forms strong oxidative derivatives including the hydroxyl radical, the nitrogen dioxide and the nitronium ion. Consequently any modification in the NO and superoxide production by the tissues can lead to the formation of strong secondary oxidative radicals (Deliconstantinos, G., Villiotou, V., Stavrides, J.C., Cancer Mol. Biol. 1: 77-86, 1994). Finally ONOO⁻ and its esters (RO-ONO or RO-ONO₂) tend to cause inactivation of the alpha -1-proteinase inhibitor (a1PI). This can be justified by the facts that: a) the hydrogen peroxide alone does not cause quick inactivation of the a1PI but acts only in

the presence of NO whereupon ONOO⁻ is formed and quick inactivation of the a1PI occur, b) solutions of tert-boutyl peroxyntirite (RO-O-O-NO₂) or ONOO⁻ cause inactivation of a1PI by themselves, and c) amines and amino acids protect the a1PI proteinase from quick inactivation (Moreno, J.J., and Pryor, W.A., Chem. Res. Toxicol. 5: 425-431, 1992). Apart from the free radicals contained in the cigarette smoke the activated alveolar macrophages represent another important source of free radical production by smokers. The alveolar macrophages activated by cigarette smoke undergo a respiration burst resulting in increased production of oxygen free radicals (mainly O₂⁻, NO and H₂O₂). Smokers appear to have an increased number of both alveolar macrophages and circulating neutrophils. The oxygen free radicals of the cigarette smoke have also been implicated in the development of lung cancer. The inhaled cigarette smoke causes increased oxidative stress in the lung cells resulting in the reduction in the concentration of the intracellular antioxidants. H₂O₂ reacts, through the production of hydroxyl radicals, with the DNA of the cells and causes a break in the double strand. As this break can be prevented by the addition of catalase, this indirectly confirms the damaging effects of H₂O₂ and the hydroxyl radicals on cellular DNA (Leanderson, P., Ann. N.Y. Acad. Sci. 686: 249-261, 1993). Furthermore H₂O₂ can cause transformation in the tracheal epithelium of the lung and has been linked to the development of bronchogenic carcinoma in smokers. Thus the detrimental role of H₂O₂ (contained in the cigarette smoke) in the lung cells and in the development of lung cancer is strongly suggested. The tar from cigarette smoke contains both semiquinone radicals and iron thus creating a system for hydroxyl radical production. The various trace elements contained in the tar of the cigarette smoke (Fe, Cu, Mn, Cd) can act both intracellularly and extracellularly. The Fe²⁺ with the well known Fenton reaction:

$$\text{Fe}^{2+} + \text{H}_2\text{O}_2 \longrightarrow \text{Fe}^{3+} + \text{OH}^\cdot + \text{OH}^-$$

causes a plethora of oxidative reactions through hydroxyl radicals. Similar production of hydroxyl radical can be achieved by Cd²⁺. Mn²⁺ is a characteristic stimulator of soluble guanylate cyclase activity. Cd²⁺ contained in the cigarette smoke is exceptionally toxic to the lung. Smokers appear to have twice the normal concentration of Cd²⁺ in their lungs. It is suggested that Cd²⁺ displaces Zn²⁺ in presentation of normalcy in the endothelium of the lung vessels (Kostial, K., In: "Trace Elements in Human and Animal Nutrition" (ed. W. Mertz) Fifth edit. Vol. 2: 319-345, Academic Press, Inc. Orlando, FL., 1986). Aldehydes, present in the cigarette smoke, react with the -SH and

-NH₂ groups of the proteins ultimately to become inert. Crotonaldehyde (α , β unsaturated aldehyde contained in cigarette smoke decreases the concentration of the -SH groups and increases the concentration of the carbonyl proteins (Stadtman, E.R., Science 257: 1220-1224, 1991).

5 Today filters on cigarettes are strongly recommended. The ultimate aim in adding filters to the cigarette is to achieve the maximum retention of noxious compounds present both in the gas and solid phases of the cigarette smoke. Epidemiological studies in smokers have shown that there was a dose-dependent response regardless of whether the cigarette smoke was administered in the gas phase, the solid phase or
10 the solid phase or the combined phase (Surgeon General of the U.S. Public Health Service. The health consequences of using smokeless tobacco, N.H. Publ. No 86-2874, Bethesda, MD, 1986). It was proven that modification of the cigarette is in itself a practical approach to reducing the noxious compounds contained in cigarette smoke. This was initially achieved using common filters and then by changing the composition
15 of the tobacco through chemical processing. Changes in the manufacturing of the cigarettes were also made with the use of porous paper or paper made of tobacco leaves. In the last 15 years many attempts have been made to make smoking less damaging to ones health by: reducing the quantity of the smoke per cigarette: changing the diameter of the cigarette; and by using perforated filters. Perforated filters allow
20 for the dilution of cigarette smoke with air to up to 50%. Activated charcoal has also been used in combination with perforated filters. This has contributed to drastic reduction in smoke yields of tar and nicotine. Such techniques are being used particularly in the developed countries like Austria, Canada, France, Germany, Sweden, England and the U.S.A.. The average yield of tar and nicotine in an American cigarette was reduced
25 from 38 mg and 2.7 mg in 1955 to 13 mg and 1 mg in 1991 respectively. In the European Community this trend towards reduction in the yields of tar and nicotine in cigarette smoke is still being continued. The upper allowable limit for tar as of Jan 1993 is 15 mg which is to be reduced to 12 mg by the beginning of Jan 1998. However in other countries the yield of tar in cigarette smoke is at 22 mg (Mitacek, E.J., Brun
30 neman, K.D., Pollednak, A.P., Hoffman, D., and Suttajit, M., Prev. Med. 20: 764-773, 1991). The changes made in the manufacturing of cigarettes led to the specific removal of certain toxic substances from the cigarette smoke; more specifically the cellulose acetate filters were introduced thus allowing for the partial removal of the semivolatile

phenols and the volatile N-nitrosamines (Brunnemann, K.D., Hoffman, d., Recent. Adv. Tobacco Res. 17: 71-112, 1989). Carbon monoxide is selectively reduced with the use of perforated filters. The concentration of carcinogenic polynuclear aromatic hydrocarbons (PAH) was selectively reduced with the use of tobacco enriched with nitrite.

- 5 However the reduction of PAH in tobacco using high concentrations of nitrite led to undesirable increases of carcinogenic N-nitrosamines, it was thus necessary to reduce the PAH by alternate means (Hoffman, D., Hoffman, I., Wynder, E.1., Lung Cancer and the Changing Cigarette in Relevance to Human Cancer of N-Nitroso-compounds, Tobacco Smoke and Mycotoxins. (eds. O'Neil, I.K., Chen, J., and Bartsch, H.) Vol. 105: 449-459, 10 1991).

From the above mentioned it becomes clear that there is a necessity to manufacture a filter capable of withholding the noxious nitrogen oxides, the free radicals, the hydrogen peroxide, the aldehydes, and the carcinogenic nitrosocompounds which are all responsible for the damaging effects of cigarette smoke on the respiratory and cardiovascular 15 systems. For the identification of the noxious compounds contained in the cigarette smoke we have conducted chemical, biological experiments. The chemical experiments performed are the following:

- a) Identification and quantitative determination of NO and NO_x using a novel chemical and biological method (this method was developed in our lab).
- 20 b) Identification of the free radicals using the lucigenine-dependent chemiluminescence methods.
- c) Identification of the aldehydes and quinone through stimulation of the enzymatic system luciferine-luciferase (this method was also developed in our lab).
- d) Identification and quantitative determination of the trace elements using the 25 method of the oxidation of luciferine by luciferase in the presence of ATP (this method was developed in our lab).
- e) Identification and quantitative determination of H₂O₂ using the isoluminol-microperoxidase dependent chemiluminescence method.
- f) Identification and quantitative determination of ONOO⁻ spectrophotometrically 30 and by luminol enhanced chemiluminescence method.
- g) Identification of the carcinogenic nitroso compound by luminol enhanced chemiluminescence.

The Biological experiments performed are the following:

- a) Identification of NO by using isolated soluble guanylate cyclase activity as functional parameter.
- b) Identification of ONOO^- by using the estimation of the oxidative stress of the human erythrocytes induced by ONOO^- .
- c) Identification of CO by using isolated soluble guanylate cyclase activity as functional parameter.

Furthermore we performed the following in vitro experiments:

- a) Isolation of alveolar macrophages from rat lung.
- b) Estimation of the oxidative stress of alveolar macrophages induced by tert-butyl-hydroperoxide (t-BHP).
- c) Determination of $\text{NO}/\text{NO}_2^-/\text{ONOO}^-$ produced by alveolar macrophages.
- d) Determination of H_2O_2 produced by alveolar macrophages.
- e) Effect of exogenous H_2O_2 on NO production by alveolar macrophages.

15

Experiments in vivo in human volunteers were performed for the determination of the following compounds:

- a) Determination of NO in the exhaled air of non-smokers.
- b) Determination of NO in the exhaled air of smokers.
- c) Determination of NO in the exhaled cigarette smoke.
- d) Determination of ONOO^- in the exhaled cigarette smoke.
- e) Determination of free radicals in the exhaled cigarette smoke.
- f) Determination of aldehydes in the exhaled cigarette smoke.

25 For the determination of NO, NOx contained a) in cigarette smoke, b) released by alveolar macrophages after challenging with cigarette smoke and c) in the exhaled cigarette smoke of human volunteers we designed and fabricated a chamber from 2.5 cm diameter, solid rods of clear Plexiglas which were hollowed out from one end with a machine-lathe to create an identical conical cavity within each of the Plexiglas rods.

30 They were then further machined and polished at the open ends, to form a mated beveled union, creating a very tight fit between the two conical cavities. A thin square of teflon sheet (polytetrafluorethylene 0.0015 inches in thickness) was sandwiched between the assemblies which were recompressed with the thumb-screws. The two tube-

access-parts at either side of the membrane, allows biologically active samples and reactive substances to be injected into, withdrawn from or modified at either side of the membrane during biological reactions (Figure 1).

5 A. Determination of NO by Chemiluminescence.

The standard NO solution was prepared according to the literature (Deliconstantinos, G., Villiotou, V., Fassitsas, C., (1992) J. Cardiovasc. Pharmacol. 12, S63-S65) and (Deliconstantinos, G., Villiotou, V., Stavrides, J.C., (1994) In: "Biology of Nitric Oxide", eds. Feelish, M., Busse, R., Moncada, S., Portland Press, in press). The reaction solution consisted of Hank's Balanced Salt Solution (HBSS) pH 7.4; H₂O₂ (500 μ M);
10 luminol (30 μ M) and the total volume was 500 μ l. The vial was vigorously stirred and the emission was recorded in Bedrthold AutoLumat LB953 luminometer.

B. Chemical Determination of NO/NO₂⁻

The chemical determination of NO was based on the diazotization of sulfanilamide by NO at acidic pH and subsequent oxidation of scopoletin which can be detected
15 fluorometrically as previously described (Deliconstantinos, G., Villiotou, V., Fassitsas, C., J. Cardiovasc. Pharmacol 12: S63-S65, 1992). Alveolar macrophages in HBSS (10⁶ cells/ml) were mixed with 100 μ l of a reagent consisting of: 20% sulfanilamide in 20% H₃PO₄ and 25 μ M scopoletin. The decay of the fluorescence was monitored at room
20 temperature (22°C) with an Aminco SPF-500 Fluorescence Spectrophotometer. The fluorescence was monitored continuously in time until the slope of the line could be measured (approx. 8 min). Slope measurements were then converted to nmols of NO using a standard curve constructed with various concentrations of pure NO. Nitrite (NO₂⁻) the end product of NO synthesis was measured on the basis of their accumulation in the supernatants of cells cultured by its reaction with Griess reagent.
25

C. Spectroscopical Determination of Peroxynitrite (ONOO⁻)


ONOO⁻ was synthesized, titrated, and stored as previously described (Deliconstantinos, G., Villiotou, V., Stavrides, J.C., In: "Biology of nitric oxide" (eds. Feelisch, M., Busse, R., and Moncada, S.) Portland Press (in press). Because of the
30 instability of ONOO⁻ at pH 7.4, UV spectra were recorded immediately after mixing the H₂O₂ and NO solution. The concentration of ONOO⁻ was determined based on an ϵ_{302} nm value of 1670 M⁻¹ cm⁻¹. UV spectra were shown after subtraction of the basal UV spectra of H₂O₂ at corresponding concentrations.

D. Estimation of free radicals

- The estimation of free radicals was performed by using the lucigenin/DAMCO (1,4 diazabicyclo-[2,2,2]octane)-induced chemiluminescence as previous described (Delicostantinos, G., Krueger, G.R.F., J. Viral Dis. 1: 22-27, 1993). The reaction mixture consisted of HBSS pH 7.4; lucigenin (30 μ M); DAMCO (100 μ M). The vial was vigorously stirred and the emission was recorded in a Bedrthold AutoLumat LB953 luminometer. Scavengers of oxygen free radicals were used (SOD, mannitol, histidine, methionine).

E. Estimation of trace elements and aldehydes

- The assays were based on the luciferase-catalyzed oxidation of D-luciferin in the presence of an ATP-magnesium salt according to the reaction:



- The trace elements Cd^{2+} , Cu^{2+} , Fe^{2+} increase the luciferase activity and the maximum chemiluminescence response is proportionally increased according to the concentrations of the trace elements up to 10 μ g. The reactions take place in HBSS pH 7.4 in total volume of 0.5 ml.

- For the estimation of the aldehydes the same enzymatic system luciferin/luciferase was used but in the absence of ATP. Aldehydes reacts with the enzymatic system to produce chemiluminescence without the presence of ATP. The reagents used were taken from an ATP assay Kit (Calbiochem-Novabiochem CA, U.S.A.).

F. Isolation of alveolar macrophages

- In brief, rats were killed with an intravenous injection of sodium pentobarbital, the thorax was opened, the lungs were perfused free of blood with Ca^{2+} free cold (4°C) phosphate buffered saline (PBS; pH 7.4), and removed intact from the chest cavity. The homogenate of rat lung was obtained by repeatedly drawing the tissue through a syringe and then passing it through successively finer stainless steel screens ranging from 32, 62 and 68 pores per inch., meshes respectively, and under a constant stream of Finkelstein Balanced Salt Solution (FBSS; pH 7.4). The final suspension of alveolar macrophages were pooled, filtered and centrifuged at 300 X g for 10 min to pellet the cells. The cell pellet, consisting of more than 98% macrophage, was washed and resuspended in Ringer's solution. Then the procedure was repeated two times. Ap-

proximately 10×10^8 macrophages were isolated per rat. Viability was assessed by trypan blue exclusion.

F. Identification of nitrosocompounds

5 Nitrosocompounds were identified by the slow release of nitric oxide (NO) after their treatment with H_2O_2 . The reaction solution consisted of dimethyl nitrosamine and/or diethyl nitrosamine (1 μ M); H_2O_2 (500 μ M); luminol (30 μ M) in HBSS pH 7.4 total volume 0.5 ml. The vial was vigorously stirred and the emission was recorded in a Bedrthold AutoLumat LB953 luminometer. Mannitol (100 mM); DMSO (100 mM) and cysteine (3.0 mM) were used to identify the formation of $ONOO^-$.

10 G. Isolation of alveolar macrophages

In brief, rats were killed with an intravenous injection of sodium pentobarbital, the thorax was opened, the lungs were perfused free of blood with Ca^{2+} free cold (4° C) phosphate buffered saline (PBS; pH 7.4), and removed intact from the chest cavity. The homogenate of rat lung was obtained by repeatedly drawing the tissue through a
15 syringe and then passing it through successively finer stainless steel screens ranging from 32, 62 and 68 pores per inch., meshes respectively, and under a constant stream of Finkelstein Balanced Salt Solution (FBSS; pH 7.4). The final suspension of alveolar macrophages were pooled, filtered and centrifuged at 300Xg for 10 min to pellet the cells. The cell pellet, consisting of more than 98% macrophage, was washed and
20 resuspended in Ringers solution. Then the procedure was repeated two times. Approximately 10×10^8 macrophages were isolated per rat. Viability was assessed by trypan blue exclusion.

H. Oxidative stress of alveolar macrophages induced by t-butyl-hydroperoxide (t-BHP)

The generation of oxygen free radicals by alveolar macrophages induced by t-BHP (2.5
25 mM) was determined by using a luminol chemiluminescence method. The chemiluminescence response was recorded in a Bedrthold AutoLumat LB953 luminometer as previous described (Deliconstantinos, G., Krueger, G.R.F., J. Viral Dis. 1, 22-27 1993).

I. Determination of hydrogen peroxide (H_2O_2)

30 An isoluminol/microperoxidase cocktail (100 mM sodium borate, 1 mM isoluminol, 0.01 mM microperoxidase in 70% water and 30% methanol at pH 8) was prepared. 50 μ l of this reagent were mixed with the isolated alveolar macrophages (10^6 cells) in HBSS in a total volume of 0.5 ml. The chemiluminescence response was converted to nmols of

H₂O₂ using a standard curve constructed with various concentrations of pure H₂O₂.

J. Preparation and Purification of soluble Guanylate cyclase (sGC) for CO estimation.

sGC from human endothelial cells was purified by GTP- agarose chromatography.

Cytosols (10 mg protein) were added to a GTP- agarose column (1.8X9 cm) pre equi

- 5 librated with 25 mM Tris-HCl buffer pH 7.6 containing 250 mM sucrose and 10 mM MnCl₂. sGC was then eluted from the column with 5 ml equilibration buffer plus 10 mM GTP.

K. Determination of Cyclic GMP

- 10 Concentrations of cGMP were determined by radioimmunoassay after acetylation of the samples with acetic anhydride (Delikonstantinos, G., and Kopeikina, L., Anticancer Res. 9: 753-760, 1989). The reaction mixture contained triethanolamine/HCl (50 mM); creatine phosphate (5 mM); MgCl₂ (3 mM); isobutylmethylxanthine (1 mM); creatine kinase (0.6 Units); GTP (1 mM); soluble guanylate cyclase (1 µg protein) in a total volume of 150 µl. The reactions were initiated by the addition of GTP and incubated for
- 15 10 min at 37° C. The incubation medium was aspirated and cGMP was extracted by the addition of ice-cold HCl (0.1 M). After 10 min, the samples were transferred to a new plate dried, and reconstituted in 5 mM sodium acetate (pH 4.75) for cGMP determination. cGMP formed was determined using a cGMP assay kit (Amersham).

20 DESCRIPTION OF THE INVENTION

The target of the present invention is to create and apply the methods in which biological substances are used that react specifically and scavenge the following:

- 25 a) NO and NOx,
b) CO,
c) H₂O₂,
d) Free radicals,
e) Aldehyde- quinones,
f) Carcinogenic nitrosocompounds,
- 30 g) Withholding the trace elements cadmium, copper, manganese, iron etc. which are inhaled during smoking.

This invention relies heavily on the notion that:

- a) There is selection of appropriate scavengers, like hemoglobin or lysates of erythrocytes or any substance which contains stereospecifically bound iron
- b) There is selection of scavengers which contain porphyrin ring with iron (e.g. protoporphyrin)
- 5 c) There is selection of scavengers which contain porphyrin ring that does not necessarily contain iron
- d) There is selection of scavengers which contain porphyrin ring complexed with other metals, e.g. Mg^{2+} , Cu^{2+}
- 10 e) A biotechnical process will be designed for the enrichment of common conventional materials which are presently used in the production of cigarette filters which will contain the above mentioned biological substances - scavengers.

The pivotal idea in this invention lies in the concept that impregnation of common conventional cigarette filters and/or filters containing activated charcoal can be enriched

15 with the biological substances, characterized by the presence of metal ions Fe^{2+} , Cu^{2+} , Mg^{2+} complexed with the porphyrin ring, as well as Fe^{2+} bound stereospecifically to protein molecules, thus allowing the noxious compounds contained in the cigarette to be withheld before the smoker inhales the cigarette smoke. This fact is the main characteristic of the present invention and consists of an undeniable innovation

20 with great feasible industrial applications.

METHODS FOR INDUSTRIAL APPLICATION

This invention was prepared in the following way in light of its applicability to industrial

25 production levels:

A solution of 1 mg/ml of hemoglobin and/or lysate of erythrocytes in phosphate buffered saline solution (PBS) with a pH of 7.4 was prepared and added to 100 mg of activated charcoal. They were incubated for 30 min at room temperature and filtered through a S&S Carl Schleicher & Schuell Co U.S.A. filter paper. The quantity of the

30 non-absorbed hemoglobin was estimated in the filtrate spectrophotometrically. The charcoal enriched with hemoglobin was left to dry at room temperature. A quantity of 200 mg of dry charcoal enriched with hemoglobin was sandwiched between two common filters so that all cigarette smoke drawn through comes into contact with the active

groups of the molecules (Fe^{2+} , Fe^{3+} , $-\text{SH}$, $-\text{NH}_2$) (Figure 2). These compatible materials are now ready to be used for the manufacturing of the new cigarette filters which we will refer to from now on as biological filters.

Alternatively hemoglobin can be replaced by biological substances characterized by the presence of metal ions Fe^{2+} , Cu^{2+} , Mg^{2+} complexed with the porphyrin ring, as well as Fe^{2+} bound stereospecifically to protein molecules, such as transferin, catalase, protoporphyrine, cytochrome C, chlorophyll.

Alternatively, a solution of 5 mg/ml of hemoglobin and/or lysate of erythrocytes in phosphate buffered saline solution (PBS) with a pH of 7.4 was prepared and scanned at 25° C using an Acta Beckman recording spectrophotometer. An absorbance peak was consistently observed at 540 nm and 575 nm (Smith, R.P., Kruszyna, H. J. Pharmacol. Exper. Ther. 191, 557-563, 1974). Common conventional cigarette filters were impregnated with these solutions and they were air dried at 25-35° C. These compatible materials are now ready to be used for the manufacturing of the new cigarette filters which we will refer to from now on as biological filters. These new biological filters ensure that the smoke which is inhaled comes completely into contact with the active groups of the hemoglobin molecules and/or lysates of the filter without changing the physical properties or the taste of the cigarette smoke. For aesthetic reasons a small part (3 mm) of a conventional filter can be adapted to the visible end of the biological filter.

Alternative industrial production methods include the following:

A solution of 5 mg/ml of protoporphyrin in buffer solution (PBS) pH 7.4 was prepared, and scanned at 25° C using an Acta Beckman recording spectrophotometer. Excitation of protoporphyrin with ultra violet light (498-408) produced an orange-red fluorescence between 620-630 nm. The conventional filters were then impregnated (soaked) with the above solution and dried with hot air (25-35° C).

Alternatively a 5 mg/ml solution of transferine in PBS pH 7.4 is scanned using the Acta Beckman recording spectrophotometer. The ferric-transferine shows a characteristic spectrum of 470 nm. The above methods for impregnating the currently used conventional filters was used.

Alternatively a 5 mg/ml solution of catalase in PBS pH 7.4 is prepared.

The above method for the preparation of the biological filter is to be followed.

Alternatively a 5 mg/ml solution of cytochrome C in PBS pH 7.4 is prepared. The above method for the preparation of the biological filter is to be followed.

Alternatively a 5 mg/ml of chlorophyll in PBS pH 7.4 is prepared. The above method for the preparation of the biological filter is to be used.

Alternatively the above mentioned biological substances are sandwiched between two common filters in solid form so that all cigarette smoke drawn through the filter comes into contact with the active groups of the molecules (Fe^{2+} , Fe^{3+} , $-\text{SH}$, $-\text{NH}_2$).

10 ANALYSIS OF THE RESULTS.

The various biological substances used to enrich the conventional filters have been shown to retain the toxic compounds (NO , CO , free radicals, H_2O_2 , aldehydes and trace elements and nitrosocompounds) from cigarette smoke in varying degrees as can be seen in the table below:

scavengers	NO %	CO %	Free radicals %	H_2O_2 %	Aldehydes %	Nitroso-compounds %	Trace elements %
Hemoglobin	90	90	90	80	90	90	95
Transferin	85	90	60	60	60	75	50
Catalase	85	90	90	90	80	80	80
Protoporphirin	85	90	70	80	70	75	80
Cytochrome C	85	80	70	80	60	60	70
Chlorophyll	15	10	40	15	10	10	80

The degree of retention of the highly damaging substances of the cigarette smoke was obtained, and the smoke of the cigarette (20 ml) filtered through a biological filter was compared with that filtered through a conventional filter (20 ml). Only 1 ml cigarette smoke drawn through the conventional filter was compared with 40 ml of cigarette smoke drawn through a biological filter. It appears that the biological filters have 40 times the capability of retaining the trace elements as compared to conventional filters.

In the following detailed experimental description representative results are shown so as to better comprehend the activity of these biological substances.

a) Identification of NO contained in cigarette smoke using the chemiluminescence method:

- 5 NO was identified using the luminol enhanced chemiluminescence method as described in the experimental section. Figures 3 and 4 illustrate a typical experiment of NO identification and estimation, as well as its scavenging after the passage of cigarette smoke through the biological filter. It appears that more than 90% of the NO is retained by the hemoglobin. The effectiveness of the biological filter is apparent in retaining and neutralizing the NO which has been implicated in toxic reactions both in lung cells and in lung fluids especially when it is involved in the formation of the strong oxidant ONOO-
- 10

b) Identification of free radicals contained in cigarette smoke using the chemiluminescence method:

- The free radicals in cigarette smoke were identified by the chemiluminescence response caused by the system lucigenine/DAMCO after its reaction with the free radicals. Figure 5 shows a characteristic peak taken within 2 seconds of the chemiluminescence response which was inhibited 100% after the passage of the cigarette smoke through a biological filter. The retention of the free radicals by the biological filters implies that there will be reduction of oxidative stress in the alveolar macrophages which is caused by conventional cigarette smoke.
- 15
- 20

c) Identification of H_2O_2 contained in cigarette smoke using the chemiluminescence method:

- H_2O_2 was estimated by the chemiluminescence response produced by the system isoluminol/microperoxidase. Figure 6 shows the characteristic peak of chemiluminescence due to the presence of H_2O_2 in cigarette smoke. In the presence of catalase (100 units/ml) the chemiluminescence response was inhibited approximately 90%. When the cigarette smoke passed through a biological filter an 80% inhibition of the chemiluminescence response was observed. The isoluminol/microperoxidase system is specific for the identification of H_2O_2 . The free radicals contained in cigarette smoke evoke a minute chemiluminescence response after their interaction with isoluminol. This minute chemiluminescence appears to be approximately 10% of the total chemiluminescence caused by H_2O_2 in the presence of free radicals since catalase inhibits the maximum chemiluminescent
- 25
- 30

response up to 90%. The retention of H_2O_2 apparently reduces both the oxidative stress and the production of NO by the alveolar macrophages.

d) Identification of trace elements and aldehydes contained in cigarette smoke using the enzymatic system luciferine/luciferase.

- 5 Trace elements contained in the cigarette smoke were identified by their capacity to stimulate the luciferase activity. Figure 7 depicts:
- 1) the chemiluminescence response caused by the oxidation of luciferine in the presence of ATP,
 - 2) the enhanced chemiluminescence response in the presence of Cd^{2+} ions (0.5 mg),
 - 10 3) the enhanced chemiluminescence response in the presence of Cu^{2+} ions (0.5 mg),
 - 4) the enhanced chemiluminescence response caused by cigarette smoke (1ml) and
 - 5) the inhibition of chemiluminescence response (with respect to that caused by the cigarette smoke) caused by 40ml cigarette smoke when passed through the biological cigarette filter. It is obvious that the chemiluminescence response caused by trace ele
 - 15 ments contained in conventional cigarette smoke are more than 40 times higher than those passed through a biological filter. The withholding of trace elements by the biological filters may have both short term and long term effects. Short term effects could entail the inhibition of redox reactions from taking place in the lung (Fe, Mn) and long term effects could entail inhibition of damages to the constituents and substances
 - 20 in the blood(Cd).

The aldehydes contained in cigarette smoke were identified and estimated using the same enzymatic system luciferine/luciferase in the absence of ATP. Aldehydes are capable of causing oxidation of luciferine. Figure 8 shows a characteristic chemiluminescence response which could last for more than an hour. This

- 25 chemiluminescence response was inhibited 100% when the cigarette smoke used had been passed through the biological filter, suggesting that the effectiveness of the biological filter to withhold the toxic aldehydes is substantial.

e) Identification of nitrosocompounds in cigarette smoke.

- The identification of nitrosocompounds contained in cigarette smoke was obtained by
- 30 estimating the slow release of NO from nitrosocompounds after their treatment with H_2O_2 . As shown in Figure 9 a peak chemiluminescence response was obtained at approximately 900 seconds. Passage of the cigarette smoke through a biological filter showed a 90% inhibition in the chemiluminescence response observed and its peak

- was taken at approximately 1200 seconds. The slow release of NO by sodium nitroprusside (SNP) after its treatment with H_2O_2 is also shown. Figure 10 shows the slow release of NO from both: the nitrosocompounds diethyl nitrosamine and dimethyl nitrosamine; and from hemoglobin enriched with nitrosocompounds from cigarette smoke treated with H_2O_2 . It is clear that the NO release by the nitrosocompounds of the cigarette smoke, which have formed adducts with hemoglobin, follow the same pattern of NO release as the nitrosocompounds diethylnitrosamine and dimethyl nitrosamine. Figure 11 shows the release of NO by the nitrosocompounds of the cigarette smoke which have formed adducts with hemoglobin after the hemoglobin-nitrosocompound adducts were irradiated with UVB ($100\text{mJ}/\text{cm}^2$) for one minute. The NO release was estimated in the presence of H_2O_2 and gave a chemiluminescence response at 1 second. The gradual rise observed in Figure 11 is due to the effect of H_2O_2 on hemoglobin (Fenton reaction).
- 5 f) Production of NO by lung macrophages:
- In vitro experiments were carried out with the help of a special chamber that was created in our lab, and which is shown in Figure 1. The teflon membrane, separating the two compartments in the chamber, is permeable to gas NO and impermeable to NO_2^- and $ONOO^-$. Unchallenged lung macrophages isolated as described in the experimental section were suspended in HBSS buffer solution (1×10^6 cells/ml) and placed in the A compartment of the chamber. In compartment B of the chamber 2.5 ml Griess reagent or sulfanilamide/scopoletin reagent is placed. The NO, released by macrophages in compartment A, diffuses through the teflon membrane into compartment B, and binds with the Griess and/or sulfamide/scopoletin reagents where it remains trapped. This indicates that lung macrophages produce gas NO. The amount of NO now present in compartment B was then determined spectrophotometrically or fluorophotometrically. The quantities of $ONOO^-$ and NO_2^- contained in compartment A of the chamber were also determined using the Griess and/or sulfanilamide/scopoletin reagents. The above experiments were repeated after challenging the macrophages with cigarette smoke before placing them in compartment A. The results, as depicted in Figure 12, show that cigarette smoke decreases the amount of NO produced whilst increasing production of $ONOO^-$ in lung macrophages, indirectly indicating the tremendous production of both NO and O_2^- which interact to form $ONOO^-$.
- 10
15
20
25
30

Repetition of the above experiments using biological filters (i.e. in which cigarette smoke was drawn through a biological filter) showed that the biological substances used, produce the same quantities of NO_2 - and ONOO- in compartment A and similar quantities of NO in compartment B as would macrophages not challenged with cigarette smoke. In this context, the components of the Griess reaction were also used to examine the kinetics of nitrosation by intermediate(s) generated during the NO/O_2 reaction in aqueous solution at physiological pH. Addition of cigarette smoke (50 ml) to a 100mM phosphate solution pH 7.4 containing 25 mM sulfanilamine and 2.5 mM N-(1-naphthyl ethylenediamine dihydrochloride (NEDD) generated an absorption at $\lambda_{\text{max}}=496 \text{ nm}$ indicative of the characteristic azo product resulting from nitration. It is worthwhile to consider the implications of the present observations vis-a-vis the expected reactivities of NO under physiological relevant conditions, where maximal concentrations of NO in the cellular microenvironment are estimated to be in the range of 0.5-10 μM . The NO concentrations are dramatically increased during cigarette smoking with detrimental effects on the lung cells.

g) Oxidative stress of lung macrophages:

The results on the effects of cigarette smoke on the oxidative stress of lung macrophages are illustrated in Figure 13. Estimations of the oxidative stress using t-BHP, showed that cigarette smoke causes twice the oxidative stress that unchallenged macrophages do. When the cigarette smoke was passed through a biological filter the oxidative stress observed was similar to that of unchallenged lung macrophages. It is thus clearly indicated the elimination of the oxidative stress induced by cigarette smoke on macrophages. The cigarette smoke is now free of the substances that cause oxidative stress on lung macrophages.

h) H_2O_2 produced by lung macrophages:

H_2O_2 produced by macrophages challenged by cigarette smoke show more than 10 times the production rate as those macrophages not challenged. The use of a biological filter show a decrease in H_2O_2 production by 90% (Figure 14) as compared to conventional filters. It is obvious that as cigarette smoke induces oxidative stress in the macrophages it increases the production of toxic H_2O_2 by these cells.

i) Reconstitution experiments:

The amount of cyclic GMP produced by the NO released by alveolar macrophages was determined using the chamber shown in Figure 1 where soluble guanylate cyclase was

placed in compartment A and alveolar macrophages were placed in compartment B. The quantities of NO produced by the macrophages were determined over a period of 50 minutes with and without cells challenged with cigarette smoke. Macrophages challenged by cigarette smoke (10 ml) released approximately ten times less the amount of NO with respect to the untreated cells thus showing 10 times less production of cyclic GMP. The above procedure was repeated using cigarette smoke passed through a biological filter. It was shown a non statistically significant difference with respect to unchallenged macrophages (control) (Figure 15). The accumulation of NO in compartment B was increased more than 5 times when the alveolar macrophages were treated with H_2O_2 (5 mM) Figure 16. This suggests that H_2O_2 increases the production of NO by a positive feedback mechanism. The L-arginine/NO pathway in macrophages is consistent with the concept that cigarette smoke causes the release of NO/ONOO-.

k) Identification of carbon monoxide (CO) in cigarette smoke:

CO presence in cigarette smoke was determined using the biological method based on the stimulation of soluble guanylate cyclase by CO. Introduction of HBSS saturated with cigarette smoke into compartment A of the chamber, in the presence of superoxide so as to neutralize NO, and the introduction of soluble guanylate cyclase into compartment B resulted in an increase in cyclic GMP production due to CO diffusing from compartment A to compartment B. Passage of cigarette smoke through a biological filter reduces the amount of cyclic GMP produced by approximately 80% (Figure 17). The above data indicates that the noxious substances NOx and CO contained in cigarette smoke are retained and neutralized by the biological filters.

25 IN VIVO EXPERIMENTS

a) We first confirmed the presence of NO and ONOO- in exhaled cigarette smoke. Human volunteers smoking a cigarette bearing a conventional filter NO present in the exhaled cigarette smoke was identified after the introduction of the exhaled smoke into an acid solution(50ml) pH 4. NO concentration was estimated by the luminol enhanced chemiluminescence method described in the experimental section, using standard curves made by commercial NO. NO concentration was found to be 0.045 mM. The experiments were repeated using biological filters and the NO concentration in the inhaled smoke was approximately 70% lower compared with the conventional filter

(Figure 18). Concentration of ONOO⁻ was determined using a solution of NaOH 1.2M which showed an increase in absorption at 303 nm (Figure 19) ($\epsilon_{303nm} = 1670 \text{ M}^{-1}\text{cm}^{-1}$). Our experiments showed that during smoking the exhaled smoke contains large quantities of ONOO⁻ (passage of 50ml exhaled smoke into 5ml NaOH 1.2M yielded a solution of 0.9 mM ONOO⁻). The ratio of NO/ONOO⁻ in the exhaled smoke was determined to be 1:20.

Therefore it appears that NOx in the lung is transformed to ONOO⁻ when it reacts with superoxide in the lung. Superoxide is released from both macrophages and redox reactions occurring in the lung during smoking. Cigarette smoke drawn by a pump does not contain ONOO⁻, however a quantity of NOx reacts with superoxide or oxygen to form nitrite ions (NO₂⁻). ONOO⁻ is formed only when cigarette smoke enters the lungs. The use of biological filters reduces the exhaled quantities of NO and ONOO⁻ by 70%.

b) ONOO⁻ reacts with bicarbonate ions of the human erythrocytes according to the reaction



The bicarbonate radical oxidizes luminol as well as aromatic and heterocyclic molecules. Alternatively ONOO⁻ may peroxidize bicarbonate to peroxybicarbonate another strong oxidizing species. On the other hand superoxide dismutase (SOD) catalyzes the nitration by ONOO⁻ and a wide range of phenolics including tyrosine in proteins.

Thus there are several potential mechanisms by which bicarbonate and SOD could influence the overall reactivity of ONOO⁻ in the cells. The presence of ONOO⁻ formed in the lungs by inhaled cigarette smoke, exhibits a dramatic increase in the oxidative stress in erythrocytes which was detected by a chemiluminescence response occurring within 5 seconds. The same experiment conducted using a biological filter resulted in an almost 100% inhibition of oxidative stress in human erythrocytes (Figure 20). Hemoglobin or erythrocyte lysates exposed to ONOO⁻ (contained in the exhaled cigarette smoke) caused the abolition of the two peaks at 540 and 575 nm normally observed in hemoglobin. The results of a representative experiment similar to the one described above was performed in 12 volunteers and is shown in Figure 21. When hemoglobin and/or lysate were exposed to a small quantity of exhaled smoke (10ml) a shift of the peaks from 540 and 575 to 525 and 555 nm was observed consistent with the formation of nitrosyl hemoglobin. The experiments were repeated using biological

filters. The peaks observed maintained their characteristic wavelengths.

- d) Aldehydes were identified in the exhaled cigarette smoke from human volunteers by their characteristic chemiluminescence peak. The experiments were repeated using biological filters and a 90% reduction of the chemiluminescent response, was observed as compared to a maximum chemiluminescence response observed when using a conventional filter (Figure 22). It is obvious that the biological filters withhold and neutralize the aldehydes in cigarette smoke whilst retaining the oxidants, thus apparently inhibiting the initiation of redox reactions from taking place in the lung which would result in the production of endogenous aldehydes.
- e) Free radicals were identified in the exhaled cigarette smoke, from human volunteers by their characteristic chemiluminescence peak. Human volunteers used cigarettes bearing conventional and biological filters. They were advised to exhale cigarette smoke (50ml) in an acid solution (0.01 N HCl) (50ml) pH: 6 and the chemiluminescence response was taken after 5 min and 60 min. At pH: 6 the exhaled ONOO⁻ is spontaneously decomposed. Within 5 min there was a 160% increase of the chemiluminescence response in the exhaled smoke passed through a conventional filter as compared to cigarette smoke passed through a biological filter (Figure 23). When the saturated by the exhaled smoke acid solution was left for an hour the difference in the chemiluminescence response increased from 160% to 250% (Figure 24). This is consistent with the concept that redox reactions are taking place continuously in the cigarette smoke through the quinone radicals and produce a series of activated oxygen species that can cause biological damage.

COMMENTARY

- Our studies have shown that alveolar macrophages possess an endogenous NO synthase, like other cells, and are capable of releasing NO/ONOO⁻ for prolonged time periods following exposure to cigarette smoke. Furthermore, once NO begins to be released by these cells, the production of NO becomes self supporting even after the stimulus is removed. Such a reaction accounts for the ability of the cigarette smoke derived NO to stimulate alveolar macrophages in releasing NO and ONOO⁻ for a period of several hours after the removal of the stimulus. Such a reaction may be initiated by the production of H₂O₂ in the lungs upon stimulation of alveolar macrophages by cigarette smoke. H₂O₂ may stimulate NO synthase activity of the lung cells to produce

- NO and ONOO- for a time period of more than an hour after the removal of the stimuli. Our experiments indeed showed that passage of cigarette smoke through a biological filter resulted in a 90% reduction (as compared to a conventional filter) of the oxidative stress in the rat alveolar macrophages. An ONOO- radical formed in the lungs may
- 5 possibly attack and inactivate the α 1-proteinase inhibitor (α 1PI). Inhibition of the α 1PI in human lungs often causes emphysema in which lung capacity is reduced. Statistical evidence indicates that smoking predisposes one to the development of emphysema (Southon, P.A., Pwis, G., Free Radicals in Medicine. Involvement in human Disease. Mayo Clin. Proc. 63: 390-408, 1988). In in vivo experiments performed in 12 volunteers
- 10 smokers a 90% reduction of the exhaled NO/ONOO- was shown when the inhaled cigarette smoke was passed through a biological filter.
- Oxygen free radicals have also been implicated in the pathogenesis of IgA immune complex induced alveolitis. Pretreatment of animals with superoxide dismutase, catalase, the iron chelator desferioxamine, or the hydroxyl radical scavenger DMSO,
- 15 supresses the development of lung injury. In contrast, the lungs of untreated positive control animals are characterized by the presence of increased numbers of alveolar macrophages. Interstitial edema and hemorrhage are also present. Furthermore, in this model of lung injury, the L-arginine is also highly protective as demonstrated by reduced: vascular permeability; vascular hemorrhage; and injury to vascular endothelial
- 20 and alveolar epithelial cells. These findings suggest that the macrophages are the source of the damage causing NO, O₂·, H₂O₂ and OH compounds (Mulligan, M.S., Jonhson, K.J., Ward, P.A., In: "Biological Oxidants: Generation and Injurious Consequences" (eds. Cochrane, C.G., and Gilbrone, M.A., Jr. Academic Press 157-172, 1992).
- 25 The retention and neutralization of the oxidants contained in the cigarette smoke by the biological filters may play a significant role in reducing the activity of the redox enzymes which are directly related to the oxidative stress in the lung cells. Biological filters drastically reduce the oxidative stress caused by inhaled cigarette smoke. Oxidative stress in the lung macrophages and endothelial cells of the lung vessels may be induced by NO,
- 30 NOx oxygen radicals and/or aldehydes contained in the cigarette smoke. Furthermore the retention of aldehydes and trace elements (especially of Cd) by the biological filters may have considerable long term effects in preserving the plasma antioxidants and in inhibiting the development of artherosclerosis. Hemoglobin contains several

neutrophilic centers which undergo covalent reactions with electrophiles. These centers induce the N-terminal valine residues of the α - and β - chain, the N1 and N3 atoms of histidine residues and the sulphhydryl group of cysteine residues. The carcinogenic nitrosocompound 4- (methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) present in

5 tobacco is transferred to the smoke during burning of cigarette and its levels in mainstream smoke could vary from 4 to 1700 ng per cigarette NNK can form adducts with hemoglobin (Hecht, S.S., Karan, S., and Carmella, S.G., in: "Human carcinogen

10 tobacco chewing and smoking. However, the statistics on current smokers, indicate that a strong case can be made for the need to reduce exposure to tobacco carcinogens and to modify their mode of action. Principal approaches toward this goal are: 1) modification of tobacco products, 2) inhibition of the metabolic activation of tobacco

15 nutrients and chemopreventing agents and 3) retention of tobacco carcinogens using specific filters which will be adapted in the tobacco of the cigarettes. Our invention using biological substances for the manufacturing of biological filters finally concerns the discovery that nitrosocompounds present in the inhaled cigarette smoke are with-

20 the non-smokers as well.

CLAIMS

1. A methodology was developed for the retention and neutralization of noxious compounds in cigarette smoke (NO, NO_x, free radicals, aldehydes, H₂O₂, CO, trace elements, and carcinogenic nitrosocompounds) which are insufficiently retained by conventional cigarette filters. This methodology is characterized by the enrichment of common, conventional cigarette filters made of fiber matrix or activated charcoal with biological substances containing iron, copper, and/or magnesium, complexed with porphyrin ring and stereospecifically bound iron in protein molecules either separately or in combinations. The enrichment of the conventional filter or activated charcoal with the above substances does not alter either the physical properties of the cigarette smoke (odor, taste and appearance) or the physical properties of the filter itself.
2. A method in accordance with claim 1, is characterized by the manufacture of a solution of hemoglobin and/or lysate 5-10 mg/ml, (quantity indicative depending upon the quality of the tobacco and its wrapping), prepared in phosphate buffer (PBS) with a pH of 7.4. Common conventional cigarette filters are immersed into the above biological solution. Then the impregnated biological filters are air dried at 25-35° C. This method ensures at least, 90% retention of NO, 90% retention of CO, 90% retention of free radicals, 90% retention of aldehydes, 90% retention of carcinogenic nitrosocompounds, 80% retention of H₂O₂ and 95% retention of trace elements.
3. A method in accordance with claims 1 and 2, is characterized by the use of hemoglobin and/or lysate in solid form. Quantities of 5-10 mg hemoglobin and/or lysate (quantity indicative depending upon the quality of tobacco and its wrapping) are sandwiched between two parts of a common, conventional cigarette filter. This method ensures the results stated in claim 2 during smoking.
4. A method in accordance with claims 1 and 2, is characterized by the preparation of a solution of 1 mg/ml (indicative only) of hemoglobin and/or lysate of erythrocytes in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal, are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with hemoglobin is sandwiched between two parts of a common, conventional filter so that all cigarette smoke drawn through the filter comes into contact with the active groups of the molecules (Fe²⁺, Fe³⁺, -SH, -NH₂). This method also ensures the results stated in claim 2, during smoking.

5. A method in accordance with claim 1, is characterized by the manufacture of a solution of transferin 5-10 mg/ml, (quantity indicative depending upon the quality of the tobaccos and its wrapping) prepared in phosphate buffer (PBS) with a pH of 7.4. Common, conventional cigarette filters are immersed into the above biological solution.
- 5 Then the impregnated biological filters are hot air dried at 25-35° C. This method ensures at least, 85% retention of NO, 90% retention of CO, 60% retention of free radicals, 60% retention of aldehydes, 75% retention of carcinogenic nitrosocompounds, 60% retention of H₂O₂ and 50% retention of trace elements.
6. A method in accordance with claims 1 and 5, is characterized by the use of transferin in solid form. Quantities of 5-10 mg transferin (quantity indicative depending upon the quality of the tobacco and its wrapping) are sandwiched between two parts of common, conventional cigarette filter. This method ensures the results stated in claim 5 during smoking.
- 10 7. A method in accordance with claims 1 and 5, is characterized by the preparation of a solution of 1 mg/ml (indicative only) of transferin in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with transferin is sandwiched between two parts of common conventional filter so that all cigarette
- 15 smoke drawn through the filter comes in to contact with the active groups of the molecules (Fe²⁺, Fe³⁺, -SH, -NH₂). This method ensures the results stated in claim 5 during smoking.
- 20 8. A method in accordance with claim 1, is characterized by the manufacture of a solution of catalase 5-10 mg/ml, (quantity indicative depending upon the quality of the tobacco and its wrapping) prepared in phosphate buffer (PBS) with a pH of 7.4. Common conventional cigarette filters are immersed in the above biological solution. Then the impregnated biological filters are air dried at 25-35° C. This method ensures at least, 85% retention of NO, 90% retention of CO, 90% retention of free radicals, 80% retention of aldehydes, 80% retention of carcinogenic nitrosocompounds, 90% retention of H₂O₂ and 80% retention of trace elements.
- 25 30 9. A method in accordance with claims 1 and 8, is characterized by the use of catalase in solid form. Quantities of 5-10 mg catalase (quality indicative depending upon the quality of tobacco and its wrapping) are sandwiched between two parts of common

conventional cigarette filter. This method ensures the results stated in claim 8 during smoking.

10. A method in accordance with claims 1 and 8, is characterized by the preparation of a solution of 1 mg/ml (indicative only) of catalase in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with catalase is sandwiched between two parts of common conventional filter so that all cigarette smoke drawn through the filter comes in to contact with the active groups of the molecules (Fe^{2+} , Fe^{3+} , -SH, - NH_2). This method ensures the results stated in claim 8 during smoking.

11. A method in accordance with claim 1, is characterized by the manufacture of a solution of protoporphyrin 5-10 mg/ml, (quantity indicative depending upon the quality of the tobacco and its wrapping) prepared in phosphate buffer (PBS) with a pH of 7.4.
- 15 Common, conventional cigarette filters are immersed in the above biological solution. Then the impregnated biological filters are air dried at 25-35° C. This method ensures at least, 85% retention of NO, 90% retention of CO, 70% retention of free radicals, 70% retention of aldehydes, 75% retention of carcinogenic nitrosocompounds, 80% retention of H_2O_2 and 80% retention of trace elements.

- 20 12. A method in accordance with claims 1 and 11, is characterized by the use of protoporphyrin in solid form. Quantities of 5-10 mg of protoporphyrin (quantity indicative depending upon the quality of the tobacco and its wrapping) are sandwiched between two parts of a common, conventional cigarette filter. This method ensures the results stated in claim 11 during smoking.

- 25 13. A method in accordance with claims 1 and 11, is characterized by the preparation of a solution of 1 mg/ml (indicative only) of protoporphyrin in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with protoporphyrin is sandwiched between two parts of common conventional filter so that all cigarette smoke drawn through the filter comes in to contact with the active groups of the molecules (Fe^{2+} , Fe^{3+} , -SH, - NH_2). This method ensures the results stated in claim 11 during smoking.
- 30

14. A method in accordance with claim 1, is characterized by the manufacture of a solution of cytochrome 5-10 mg/ml, (quantity indicative depending upon the quality of the tobacco and its wrapping) prepared in phosphate buffer (PBS) with a pH of 7.4. Common conventional cigarette filters are immersed in the above biological solution.
- 5 Then the impregnated biological filters are air dried at 25-35° C. This method ensures at least, 85% retention of NO, 80% retention of CO, 70% retention of free radicals, 60% retention of aldehydes, 60% retention of carcinogenic nitrosocompounds, 80% retention of H₂O₂ and 70% retention of trace elements.
15. A method in accordance with claims 1 and 14, is characterized by the use of
- 10 cytochrome C in solid form. Quantities of 5-10 mg of cytochrome C (quantity indicative depending upon the quality of the tobacco and its wrapping) are sandwiched between two parts of a common, conventional cigarette filter. This method ensures the results stated in claim 14 during smoking.
16. A method in accordance with claims 1 and 14, is characterized by the preparation
- 15 of a solution of 1 mg/ml (indicative only) of cytochrome C in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with
- 20 cytochrome C is sandwiched between two parts of a common, conventional filter so that all cigarette smoke drawn through the filter comes into contact with the active groups of the molecules (Fe²⁺, Fe³⁺, -SH, -NH₂). This method ensures the results stated in claim 14 during smoking.
17. A method in accordance with claim 1, is characterized by the manufacture of a solution of chlorophyll 5-10 mg/ml, (quantity indicative depending upon the quality of the
- 25 tobacco and its wrapping) prepared in phosphate buffer (PBS) with a pH of 7.4. Common, conventional cigarette filters are immersed in the above biological solution. Then the impregnated biological filters are hot air dried at 25-35° C. This method ensures at least, 15% retention of NO, 10% retention of CO, 40% retention of free radicals, 10% retention of aldehydes, 10% retention of carcinogenic nitrosocompounds, 15%
- 30 retention of H₂O₂ and 80% retention of trace elements.
18. A method in accordance with claims 1 and 17, is characterized by the use of chlorophyll in solid form. Quantities of 5-10 mg chlorophyll (quantity indicative of the quality of tobacco and its wrapping) are sandwiched between two parts of a common,

conventional cigarette filter. This method ensures the results stated in claim 17 during smoking.

19. A method in accordance with claims 1 and 17, is characterized by the preparation of a solution of 1 mg/ml (indicative only) of chlorophyll in phosphate buffered saline solution (PBS) with a pH of 7.4. 100 mg (indicative only) of activated charcoal are added to this solution and the mixture is then incubated for approximately 30 min at room temperature. A quantity of 200 mg (indicative only) of dry charcoal enriched with chlorophyll is sandwiched between two parts of a common, conventional filter so that all cigarette smoke drawn through the filter comes into contact with the active groups of the molecules (Fe^{2+} , Fe^{3+} , -SH, $-\text{NH}_2$). This method ensures the results stated in claim 17 during smoking.

20. The methodology in accordance with claim 1, is characterized by the preparation of solutions and/or solid biological substances whose common features are:

- a) Any molecule containing heme iron (heme, hematin etc.).
- 15 b) Any macromolecule containing stereospecifically bound iron or copper (ferritine, ceruloplasmine, etc.).
- c) Any macromolecule containing porphirin ring which does not necessarily contain iron.
- d) Any macromolecule containing porphirin ring complexed with other metals other than iron (Mg, Cu).

21. The biochemical-pathophysiological mechanisms implicated in lung and blood (constituent and substance) damage which the present invention counters and the know-how leading to technological countermeasures. These mechanisms specifically involve the alveolar macrophages and the endothelial cells of the lung which upon being challenged by cigarette smoke undergo oxidative stress resulting in the manyfold production of NO/ONOO- and H_2O_2 for prolonged periods of time. H_2O_2 further stimulates the production of NO/ONOO- creating a vicious cycle. ONOO- radicals inhibit the α_1 -proteinase Inhibitor ($\alpha_1\text{PI}$), a significant protective mechanism in lung tissue. Also the manufacturing processes in accordance with claim 1 allowing retention and neutralization of the noxious compounds contained in cigarette smoke ie., NO, CO, aldehydes, free radicals, H_2O_2 , trace elements, carcinogenic nitrosocompounds, thus offering considerable protection both to the active and passive smoker from diseases such as lung emphysema, lung cancer, chronic bronchitis and long term diseases of

the cardiovascular system.

22. A new cigarette filter is characterized either by its enrichment with the biological substances mentioned in claims 1 and 20, or by the presence of activated charcoal enriched with the biological substances mentioned in claims 1 and 20 placed between
- 5 the two ends or at one of the two ends of the new cigarette filter. The two ends or the one end respectively of this cigarette filter are made from fiber matrix used for the manufacture of common, conventional filters.

AMENDED CLAIMS

[received by the International Bureau on 27 October 1995 (27.10.95);
original claims 1-22 replaced by amended claims 1-14 (2 pages)]

1. A filter for filtering tobacco smoke characterised in that it comprises a fiber matrix enriched with a biological substance selected from one or more substances containing iron, copper, and/or magnesium complexed with a porphyrin ring and iron bound stereospecifically in protein molecules.
2. A filter as claimed in Claim 1, characterised in that it comprises activated charcoal enriched with the biological substance.
3. A filter as claimed in Claim 1 or 2, characterised in that said enriched fiber matrix is flanked by fiber matrix which is not enriched with said biological substance.
4. A filter as claimed in any one of Claims 1 to 3, characterised in that the biological substance comprises hemoglobin and/or lysate of erythrocytes.
5. A filter as claimed in any one of Claims 1 to 3, characterised in that the biological substances are selected from iron Fe^{2+} ions bound stereospecifically to one or more of transferrin, catalase, protoporphyrin, cytochrome C and chlorophyll.
6. A filter as claimed in any one of Claims 1 to 5, wherein the biological substance is in solid form.
7. A cigarette characterised in that it is provided with a filter as claimed in any one of Claims 1 to 6.
8. A method of making a tobacco smoke filter as claimed in any one of Claims 1 to 6, comprising impregnating a conventional tobacco smoke filter with one or more of said biological substances.
9. A method as claimed in Claim 8, characterised in that the filter comprises activated charcoal.
10. A method as claimed in Claim 8 or 9, characterised in that the biological substance comprises hemoglobin and/or lysate of erythrocytes.
11. A method as claimed in any one of Claims 8 to 10, characterised in that the biological substance is provided as a 1-10 mg/ml solution in a phosphate buffered saline solution having a pH of 7.4.
12. A method of filtering tobacco smoke comprising providing a filter as claimed in any one of Claims 1 to 6 and passing tobacco smoke therethrough.

- 5 13. A method as claimed in Claim 12, characterised in that the filter retains from 15 to 90% NO ; 10 to 90% CO ; 40 to 90% free radicals; 10 to 90% aldehydes ; 10 to 90% carcinogenic nitroso compounds; 15 to 90% H₂O₂ ; and 50 to 95% of trace elements present in the tobacco smoke before passing through the filter.
- 10 14. A method as claimed in Claim 13, characterised in that the filter retains from 85 to 90% NO ; 80 to 90% CO; 60 to 90% free radicals; 60 to 90% H₂O₂ ; 60 to 90% aldehydes ; 60 to 90% carcinogenic nitroso compounds ; and 70 to 95% of the trace elements present in the tobacco smoke before passing through the filter.




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5




Figure 6




Figure 8.

(A) the slow release of NO from the nitrosocompounds after their treatment with H_2O_2 is shown.




Figure 9.




Figure 10.




Figure 11.

11/24

SUBSTITUTE SHEET (RULE 26)




Figure 12.




Figure 13.




Figure 14.




Figure 15.




Figure 16.




Figure 17.




Figure 18.




Figure 19.




Figure 20.

20/24

SUBSTITUTE SHEET (RULE 26)




Figure 21.




Figure 22.




Figure 23.




Figure 24.

24/24

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GR 94/00015

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A24D3/14 A24D3/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A24D A23D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB,A,2 083 998 (ADVANCE KK) 7 April 1982 see the whole document ---	1-3,8,9, 11,12, 20,21
X	US,A,4 049 673 (I.H. SCHEINBERG) 20 September 1977 see column 7, line 29 - column 8, line 29; claims ---	1,2,20, 21
X	CH,A,609 217 (S. NEUKOMM) 28 February 1979 see the whole document ---	1,3,20, 21
X	US,A,4 612 333 (N.I. VASSILEFF) 16 September 1986 see column 10, line 4 - line 13; example 28 ---	1,3,18, 20,21
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 February 1995

Date of mailing of the international search report

09.03.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lepretre, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GR 94/00015

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 82-84398E & JP,A,57 138 375 (KOWA KK) 26 August 1982 see abstract ---	1,17,20, 21
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 88-288470 & JP,A,63 209 718 (UBE INDUSTRIES LTD) 31 August 1988 see abstract ---	1,10, 20-22
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 83-726087 & JP,A,58 107 166 (ADVANCE KAIHATSU) see abstract ---	1,20,21
X	PATENT ABSTRACTS OF JAPAN vol. 14, no. 118 (C-0697) 6 March 1990 & JP,A,01 317 538 (ASAHI CHEM. IND. CO. LTD.) 22 December 1989 see abstract -----	1,20,21

INTERNATIONAL SEARCH REPORT

Int. onal Application No
PCT/GR 94/00015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2083998	07-04-82	JP-C- 1511327	09-08-89
		JP-A- 57039767	05-03-82
		JP-B- 63057032	10-11-88
		CA-A- 1176941	30-10-84
		DE-A- 3133169	08-04-82
		FR-A, B 2490461	26-03-82
		US-A- 4414988	15-11-83

US-A-4049673	20-09-77	US-A- 3982897	28-09-76
		CA-A- 1002037	21-12-76
		DE-A- 2227787	25-01-73
		FR-A, B 2110329	02-06-72
		GB-A- 1367366	18-09-74

CH-A-609217	28-02-79	NONE	

US-A-4612333	16-09-86	US-A- 4804688	14-02-89
		US-A- 4724242	09-02-88
